

Weihmüller, M. P., G. R. Cattáneo, A. Sharpe, M. Takigami & A. D. Izeta (2023) Zooarqueología e isótopos estables en el valle de Ongamira (Depto. Ischilín, Córdoba, Argentina). Poster presentado en el IV Taller de Isótopos Estables y Arqueología en el Sur de Sudamérica, Ushuaia, abril 2023

Referencias bibliográficas

Ambrose, S. H. (1990). Preparation and characterization of bone and tooth collagen for isotopic analysis. *Journal of Archaeological Science*, 17(4), 431-451. [https://doi.org/10.1016/0305-4403\(90\)90007-R](https://doi.org/10.1016/0305-4403(90)90007-R)

Cattáneo, G. R., & Izeta, A. D. (Eds.). (2019). *Arqueología en Ongamira* (1 ed.). Consejo Nacional de Investigaciones Científicas y Técnicas. <https://suquia.ffyh.unc.edu.ar/handle/suquia/4511>

DeNiro, M. J. (1985). Postmortem preservation and alteration of in vivo bone collagen isotope ratios in relation to paleodietary reconstruction. *Nature*, 317(31), 806-809. <https://doi.org/10.1038/317806a0>

Fernández, F. J., Gil, A. F., Ugan, A., & Neme, G. A. (2016). Ecological conditions and isotopic diet (^{13}C and ^{15}N) of Holocene caviomorph rodents in Northern Patagonia. *Journal of Arid Environments*, 127, 44-52. <https://doi.org/10.1016/j.jaridenv.2015.10.019>

Giardina, M. A., Neme, G. A., & Gil, A. F. (2014). Rheidae Egg Human Exploitation and Stable Isotopes: Trends from West Central Argentina. *International Journal of Osteoarchaeology*, 24(2), 166-186. <https://doi.org/10.1002/oa.2346>

Gil, A. F., Ugan, A., Otaola, C., Neme, G., Giardina, M., & Menéndez, L. (2016). Variation in camelid $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in relation to geography and climate: Holocene patterns and archaeological implications in central western Argentina. *Journal of Archaeological Science*, 66, 7-20. <https://doi.org/10.1016/j.jas.2015.12.002>

Hyland, C., Scott, M. B., Routledge, J., & Szpak, P. (2022). Stable Carbon and Nitrogen Isotope Variability of Bone Collagen to Determine the Number of Isotopically Distinct Specimens. *Journal of Archaeological Method and Theory*, 29, 666-686. <https://doi.org/10.1007/s10816-021-09533-7>

Izeta, A. D., Cattáneo, R., Robledo, A. I., Takigami, M., Yoneda, M., Tokanai, F., Kato, K., & Matsuzaki, H. (2021). New radiocarbon evidence for human occupation in Central Argentina during the Middle and Late Holocene: The Ongamira valley case. *Radiocarbon*, 1-20. <https://doi.org/10.1017/RDC.2021.22>

Otaola, C., Ugan, A., & Gil, A. F. (2018). Environmental diversity and stable isotope variation in faunas: Implications for human diet reconstruction in Argentine mid-latitude deserts. *Journal of Archaeological Science: Reports*, 20, 57-71. <https://doi.org/10.1016/j.jasrep.2018.03.005>

Scheifler, N. A., Merino, M. L., Vitale, P., Kaufmann, C. A., Messineo, P. G., Álvarez, M. C., & Bocherens, H. (2020). Isotopic Ecology in Modern and Holocene Populations of Pampas Deer (*Ozotoceros bezoarticus*) from Eastern Central Argentina. Implications for Conservation Biology and Ecological Models of Hunter-gatherer Subsistence. *Environmental Archaeology*, 1-17. <https://doi.org/10.1080/14614103.2020.1846451>

Scheifler, N., Vitale, P., & Kaufmann, C. A. (2021). Primeros datos isotópicos ($\delta^{13}\text{C}_{\text{col}}$, $\delta^{15}\text{N}$) de una población moderna de guanaco en el Monte de Patagonia nororiental. Implicancias paleoecológicas y arqueológicas. *InterSecciones en Antropología*, 22(2), Article 2.
<https://doi.org/10.37176/iea.22.2.2021.633>

Sharpe, A. E., Emery, K. F., Inomata, T., Triadan, D., Kamenov, G. D., & Krigbaum, J. (2018). Earliest isotopic evidence in the Maya region for animal management and long-distance trade at the site of Ceibal, Guatemala. *Proceedings of the National Academy of Sciences*, 115(14), 3605-3610.
<https://doi.org/10.1073/pnas.1713880115>

Silva, L. C. R., Giorgis, M. A., Anand, M., Enrico, L., Pérez-Harguindeguy, N., Falczuk, V., Tieszen, L. L., & Cabido, M. (2011). Evidence of shift in C4 species range in central Argentina during the late Holocene. *Plant and Soil*, 349(1-2), 261-279. <https://doi.org/10.1007/s11104-011-0868-x>

Tessone, A., Rindel, D., Belardi, J. B., Panarello, H. O., & Goñi, R. A. (2014). $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ Variability in Modern Guanaco (*Lama guanicoe*) Assemblages in Southern Patagonia: Implications for Zooarchaeological Studies. *International Journal of Osteoarchaeology*, 24(2), 202-218.
<https://doi.org/10.1002/oa.2330>

Torrela S. A. y J. Adámoli (2006) Situación ambiental de la ecorregión del Chaco Seco. Pp. 75- 82, en: La situación ambiental Argentina 2005 (A Brown, U Martínez Ortiz, M Acerbi y J Corcuera, eds.). Fundación Vida Silvestre Argentina.
https://www.researchgate.net/profile/Sebastian-Torrella/publication/284500076_Situacion_ambiental_de_la_ecorregion_del_Chaco_Seco/links/5880d5c045851503b6edd8c4/Situacion-ambiental-de-la-ecorregion-del-Chaco-Seco.pdf

van Klinken, G. J. (1999). Bone Collagen Quality Indicators for Palaeodietary and Radiocarbon Measurements. *Journal of Archaeological Science*, 26(6), 687-695.
<https://doi.org/10.1006/jasc.1998.0385>

Weihmüller, M. P., Cattáneo, G. R., Takigami, M., & Izeta, A. D. (2022). Una primera aproximación a la composición isotópica de carbono ($\delta^{13}\text{C}_{\text{col}}$) y nitrógeno ($\delta^{15}\text{N}$) en restos de guanacos modernos y arqueológicos del noroeste de Córdoba, Argentina. *Revista del Museo de La Plata UNLP*, 7, 42R-43R.
<https://doi.org/10.24215/25456377e015R>

Yanes, Y., Izeta, A. D., Cattáneo, R., Costa, T., & Gordillo, S. (2014). Holocene (~4.5–1.7 cal. Kyr BP) paleoenvironmental conditions in central Argentina inferred from entire-shell and intra-shell stable isotope composition of terrestrial gastropods. *The Holocene*, 24(10), 1193-1205.
<https://doi.org/10.1177/0959683614540959>